




Sink toward Source

- Testable hypotheses
- Environmental diagnosis

¹Sanjeev Gupta, ¹Nick Warner, ²Melissa Rice, ³Jung-Rack Kim, ¹Kate Goddard, ³Lucille Le Corre, ³Jan-Peter Muller



Known knowns

- Near complete source-to-sink sediment routing system
- Closed basin
- Well preserved paleogeomorphology
- Spatial and temporal grain-size segregation
 - Can sample variability in sedimentary rock types
- Distinct stratigraphic variability.


What can this tell us?

- We can directly link erosion to depositional record
 - We are not guessing the source region
- Sedimentary lithofacies can be linked to palaeogeomorphology
 - We do not have to guess where we are in the system

Eberswalde has a testable depositional model.


Hypothesis: Fluvio-deltaic system prograding into a lake


- Multiple sediment 'lobes' that progressively step basinward
- Large-scale channel switching (avulsion)
- Evidence for increased frequency of channel bifurcation at distal ends of lobes



Channel Bifurcation

Mouth bar development leads to channel bifurcation






Interpretation:

- -multiple orders of bifurcation...or
- -topographically overlapping sinuous channels


CTX Channel Topography 0 m -1950 m -1520 Elevation (m) -1540 -1560 -1580 -1600 **Distance (m)** 2500 500 1000 1500 3000 4000 3500 4500 0

HiRISE Dtm



~1 km

Mossy delta, Canada




Edmonds & Slingerland, 2007



Spaghetti channels – little bifurcation



Photomorphology predicts distinct stratigraphy





-Basal sequence:

- light-toned, polygonally fractured layered material
- lower 50 m of delta
- individual layering, m to sub-m scale thicknesses

Upper Sequence

- Dunes obscure stratigraphy
- Bedding characteristics of light-toned beds similar to basal sequence
- Interpretations
 - Continuation of basal sequence, preferential aeolian cover on upper slopes
 - Intercalated coarse and fine sequences? (i.e. channel sands distal channel muds-silts)
 - ?Change in source material, un-related to delta processes






Distal Eberswalde

Consistent dip, < 1° towards crater interior

_	Layer ID	Dip	Dip_Dir	RMS Error
	1	0.48	352	0.82
	2	0.49	85	0.73
	3	0.60	84	1.67
	4	0.69	84	0.65
	5	0.44	74	1.20
	6	0.76	55	1.47
	7	0.90	72	1.07
	8	0.71	64	0.77
	9	0.85	71	1.61
	10	1.17	80	0.99





Distal Lobe 2: Photomorphology

We predict a vertical coarsening-up signature.


Fluvial channels

Delta front -> Lacustrine deposits?









Climbing ripple cross-laminae

- Dominate delta front mouthbar deposits



Rapid deceleration of jet flows in standing water lead to high sediment aggradation rates

Dr. Sarah Davies, Univ. Leicester

Fluvial distributary channel

Progradation of delta leads to migration of fluvial channel facies across the top of delta mouthbar facies

Distributary channel base: erosional relief of 10's cm


distributary channel sandstones 1 m part of shelf delta siltstone Tullig Cycle, Clare, Ireland, SJ Davies

Dr. Sarah Davies, Univ. Leicester

Cross-bedded sandstones in fluvial distributary channels



Dr. Sarah Davies, Univ. Leicester



Delta-lake hypothesis questions

- Fluvial -> delta -> lake transition ?
- If lake, shallow lake, deep lake or transitions
- Multiple cycles of sedimentary activity
- Timescales of deposition longevity of system

What could MSL tell us?

- Grain size and sedimentary structures
 - Sedimentary processes and proximality
- Sedimentary facies associations
 - High-resolution sedimentary environments
- Sedimentary geometries & architecture
 - Channel geometries -> palaeo-water discharge
- Provenance we may tap interesting rocks from Eberswalde catchment

What could MSL tell us?


- Vertical sedimentary successions
 - Relative temporal evolution of sedimentary systems
 - Cycles of deposition and erosion
 - Stratigraphic hiatus' time breaks
- Large-scale stratigraphy leads to:
 - Relative chronology
 - Possible big time breaks
- -> Can we say something about longevity of sedimentary system?

How can we reconstruct time?


- For timescales look for time gaps
- Hiatuses
- Look for sedimentary facies disjunctions
- Duricrusts, aeolian intercalations,

Breaking Walther's Law

Hypothesis: Does lake level fall drive basinward shift of deltas?



Youngest meander belt extends basinward of older deltaic seds = basinward shift of depositional systems? Test through stratigraphy.



Pots of gold

- Source area evolution: provenance and weathering records
- Surface system evolution
- Water fluxes from fluvial systems
- Climate records what supported lacustrine system

Key site for habitability

Conclusions

- Eberswalde has rich diversity of depositional sedimentary features that can be linked to palaeo-geomorphology.
- Sedimentary structures and architecture from MSL will provide stage for robust interpretations of process and environment.
- We can couple mineralogy to geologic context.
- We can understand issues of habitability because we know the context.